



2018 III 15

1000

Seat No. :

--	--	--	--	--

Time : 2½ Hours

MATHEMATICS (New Pattern)

Subject Code

H	7	5	4
---	---	---	---

Total No. of Questions : 30

(Printed Pages : 7)

Maximum Marks : 80

- INSTRUCTIONS:**
- 1) All questions are compulsory.
  - 2) The question paper consists of 30 questions divided into five Sections A, B, C, D and E.
  - 3) Section A contains 7 questions of 1 mark each which are multiple choice type questions, Section B contains 7 questions of 2 marks each, Section C contains 7 questions of 3 marks each, Section D contains 7 questions of 4 marks each and Section E contains 2 questions of 5 marks each.
  - 4) There is no overall choice in the paper. However internal choice is provided in 2 questions of 3 marks each, 2 questions of 4 marks each and 2 questions of 5 marks each. In questions with choices only one of the choices is to be attempted.
  - 5) Use of calculator is not permitted.

## SECTION – A

Question numbers 1 to 7 carry 1 mark each. In each question, four options are provided out of which one is correct. Write the correct option.

1. If  $f(x)$  is differentiable at  $x = a$ , which of the following statement may be false ?

- $f(x)$  is continuous at  $x = a$
- $\lim_{x \rightarrow a} f(x)$  exist
- $\lim_{h \rightarrow 0} \frac{f(a+h) - f(a)}{h} = \lim_{h \rightarrow 0^+} \frac{f(a+h) - f(a)}{h}$
- The second derivative of  $f(x)$  i.e.  $f''(x)$  exist at  $x = a$



2. If  $\vec{a} = \hat{i} + 2\hat{j} + 3\hat{k}$ ,  $\vec{b} = -\hat{i} + 2\hat{j} + \hat{k}$ ,  $\vec{c} = 3\hat{i} + \hat{j}$  are such that  $\vec{a} + t\vec{b}$  is right angled to  $\vec{c}$ , then the value of  $t$  will be equal to

- 5
- 4
- 6
- 2

3. If  $f(x) = \int_0^x t \sin t dt$  then  $f'(x)$  is

- $\cos x + x \sin x$
- $x \sin x$
- $x \cos x$
- $\sin x + x \cos x$

4. Find the general solution for the differential equation  $\frac{dy}{dx} = \sqrt{4 - y^2}$  ( $-2 < y < 2$ ).

- $2y = \sin(x + c)$
- $y = 2 \sin(x + c)$
- $x = 2 \sin(y + c)$
- $2x = \sin(y + c)$

5. For  $x \in \mathbb{R}$ ,  $\sin(\tan^{-1} x)$  is equal to

- $\frac{x}{2\sqrt{1+x^2}}$
- $\frac{2}{\sqrt{1+x^2}}$
- $\frac{1}{\sqrt{1+x^2}}$
- $\frac{x}{\sqrt{1+x^2}}$

6. Let  $A$  be a square matrix of order  $2 \times 2$ . Then  $|KA|$  is equal to

- $K|A|$
- $K^2|A|$
- $2|A|$
- $2K|A|$

7. Let  $f: \mathbb{R} \rightarrow \mathbb{R}$  be the function defined by  $f(x) = 2x^2 + 3x + 1$ . Using differential with  $h = 0.12$  in the usual notation, the approximate value of  $f(2.12)$  is

- 13.63
- 16.32
- 16.3488
- 16.35

## SECTION - B

Question numbers 8 to 14 carry 2 marks each.

8. Let  $*$  be a binary operation on the set of all non-zero real numbers given by  $a * b = \frac{ab}{5}$  for all  $a, b \in \mathbb{R} - \{0\}$ .

Write the identity element for the binary operation  $*$ . Also find the value of  $x$  given by  $2 * (x * 5) = 10$ .

9. If  $A$  and  $B$  are symmetric matrices, prove that  $AB - BA$  is a skew symmetric matrix.

10. If  $y = \begin{vmatrix} f(x) & g(x) & h(x) \\ l & m & n \\ a & b & c \end{vmatrix}$ , prove that  $\frac{dy}{dx} = \begin{vmatrix} f'(x) & g'(x) & h'(x) \\ l & m & n \\ a & b & c \end{vmatrix}$ .

11. Find the integral  $I = \int \frac{(x+1)(x+\log x)^2}{x} dx$ .

12. Form the differential equation representing the family of curves  $y = e^x (a \cos x + b \sin x)$  where  $a$  and  $b$  are arbitrary constants.

13. If  $A$  and  $B$  are independent events, then prove that  $P(A \cup B) = 1 - P(A') \cdot P(B')$ .

14. If  $f(x) = \frac{4x+3}{6x-4}$   $x \neq \frac{2}{3}$ , show that  $f \circ f(x) = x$  for all  $x \neq \frac{2}{3}$ .



## SECTION - C

Question numbers 15 to 21 carry 3 marks each.

15. Prove that  $\operatorname{cosec}^{-1} \operatorname{cosec}^{-1} x = \frac{\sqrt{1+x^2}}{\sqrt{2+x^2}}$ .

16. Using atleast three properties of determinants show that

$$\begin{vmatrix} a^2+1 & ab & ac \\ ab & b^2+1 & bc \\ ca & cb & c^2+1 \end{vmatrix} = 1+a^2+b^2+c^2$$

OR

Using atleast three properties of determinants, solve  $\begin{vmatrix} a+x & a-x & a-x \\ a-x & a+x & a-x \\ a-x & a-x & a+x \end{vmatrix} = 0$ .


17. Discuss the continuity of the function  $f(x)$  at  $x = \frac{\pi}{2}$  if

$$f(x) = \frac{1 - \sin^2 x}{3 \cos^2 x} \quad x < \frac{\pi}{2}$$

$$= \frac{1}{3} \quad x = \frac{\pi}{2}$$

$$= \frac{8(1 - \sin x)}{3(\pi - 2x)^2} \quad x > \frac{\pi}{2}$$

18. Prove that  $\int \sqrt{x^2+a^2} \, dx = \frac{x}{2} \sqrt{x^2+a^2} + \frac{a^2}{2} \log |x + \sqrt{x^2+a^2}| + c$ .



19. Evaluate the definite integral  $\int_1^4 (x^2 - x) dx$  as limit of sums.

20. Solve the differential equation  $(1 + e^{x/y}) dx + e^{x/y} \left(1 - \frac{x}{y}\right) dy = 0$ .

OR

Find the equation of a curve passing through the point  $(0, 2)$ , given that the sum of the coordinates of any point on the curve exceeds the magnitude of the slope of the tangent to the curve at that point by 5.

21. If  $\vec{a} = a_1\hat{i} + a_2\hat{j} + a_3\hat{k}$

$$\vec{b} = b_1\hat{i} + b_2\hat{j} + b_3\hat{k}$$

$$\vec{c} = c_1\hat{i} + c_2\hat{j} + c_3\hat{k}$$

then prove that  $[\vec{a} \ \vec{b} \ \vec{c}] = \begin{vmatrix} a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \\ c_1 & c_2 & c_3 \end{vmatrix}$ .

SECTION - D

Question numbers 22 to 28 carry 4 marks each.

22. If  $A = \begin{bmatrix} 2 & -3 & 5 \\ 3 & 2 & -4 \\ 1 & 1 & -2 \end{bmatrix}$  find  $A^{-1}$  using adjoint method. Using  $A^{-1}$  solve the system of equations

$$2x - 3y + 5z = 11$$

$$3x + 2y - 4z = -5$$

$$x + y - 2z = -3$$

OR



Find  $A^{-1}$  if  $A = \begin{bmatrix} 0 & 1 & 1 \\ 1 & 0 & 1 \\ 1 & 1 & 0 \end{bmatrix}$ . Also show that  $A^{-1} = \frac{A^2 - 3I}{2}$  where  $I$  is the identity

matrix of order 2.

23. If  $y = \sin(x^x) + x^{\tan x}$ , find  $\frac{dy}{dx}$ .
24. Find the area bounded by the parabola  $4y = 3x^2$  and the line  $2y = 3x + 12$  in the first quadrant.
25. Find the equation of the plane that contains the point  $(1, -1, 2)$  and is perpendicular to each of the planes  $2x + 3y - 2z = 5$  and  $x + 2y - 3z = 8$ .
26. Find the points on the line  $\frac{x+2}{3} = \frac{y+1}{2} = \frac{z-3}{2}$  at a distance of 5 units from the point  $P(1, 3, 3)$ .
27. Solve the following Linear Programming problem  
Minimize  $Z = 20x + 10y$   
Subject to
- $$\begin{aligned}x + 2y &\leq 40 \\3x + y &\geq 30 \\4x + 3y &\geq 60 \\x, y &\geq 0\end{aligned}$$
28. Bag I contains 3 red and 4 black balls and Bag II contains 4 red and 5 black balls. One ball is transferred from Bag I to Bag II and then a ball is drawn from Bag II. The ball so drawn is found to be red in colour. Find the probability that the transferred ball is black.

OR

Two cards are drawn with replacement from a well shuffled pack of 52 cards. Find the mean and variance of the number of kings.



SECTION - E

Question numbers 29 and 30 carry 5 marks each.

29. Find the co-ordinates of the point on the curve  $\sqrt{x} + \sqrt{y} = 4$  at which tangent is equally inclined to the axes.

OR

Show that the height of the cylinder of greatest volume which can be inscribed in a right circular cone of height  $h$  and semi vertical angle  $\alpha$  is one-third that of the

cone and the greatest volume of cylinder is  $\frac{4}{27} \pi h^3 \tan^2 \alpha$ .

30. Find the integral  $I = \int \frac{\sqrt{\cos 2x}}{\sin x} dx$ .

OR

Find the integral  $I = \int \frac{1}{\cos^6 x + \sin^6 x} dx$ .

---